

Package ‘xtfifevd’

May 8, 2026

Type Package

Title Panel Fixed Effects Filtered and Variance Decomposition Estimation

Version 1.0.2

Description Implements fixed effects estimators for time-invariant variables in panel data models. Provides three estimation methods: FEVD (Fixed Effects Vector Decomposition) from Plumper and Troeger (2007) [<doi:10.1093/pan/mpm002>](https://doi.org/10.1093/pan/mpm002), and FEF (Fixed Effects Filtered) and FEF-IV (instrumental variables variant) from Pesaran and Zhou (2018) [<doi:10.1080/07474938.2016.1222225>](https://doi.org/10.1080/07474938.2016.1222225). All methods use the correct Pesaran-Zhou variance estimators that account for generated regressor uncertainty, avoiding the size distortions documented in the literature.

License GPL-3

URL <https://github.com/muhammedalkhalaf/xtfifevd>

BugReports <https://github.com/muhammedalkhalaf/xtfifevd/issues>

Encoding UTF-8

RoxygenNote 7.3.3

Depends R (>= 4.1.0)

Imports stats, MASS

Suggests plm, lmtest, sandwich, testthat (>= 3.0.0), knitr, rmarkdown

Config/testthat/edition 3

NeedsCompilation no

Author Muhammad Alkhalaf [aut, cre, cph] (ORCID: [<https://orcid.org/0009-0002-2677-9246>](https://orcid.org/0009-0002-2677-9246))

Maintainer Muhammad Alkhalaf <muhammedalkhalaf@gmail.com>

Repository CRAN

Date/Publication 2026-04-28 20:30:02 UTC

Contents

bw_ratio	2
xtfifevd	3
Index	7

bw_ratio	<i>Between/Within Variance Ratio for Time-Invariant Variables</i>
----------	---

Description

Computes the between-panel and within-panel standard deviations for specified variables, along with their ratio. This diagnostic helps assess whether FEVD/FEF methods may improve upon standard FE estimation.

Usage

```
bw_ratio(data, variables, id)
```

Arguments

data	A data frame containing the panel data.
variables	A character vector of variable names to analyze.
id	Character string naming the panel identifier variable.

Details

For truly time-invariant variables, the within-panel SD should be zero (or near-zero due to numerical precision), giving an infinite ratio.

According to Plümper and Troeger (2007), FEVD/FEF methods tend to improve upon standard FE when:

- The between/within ratio exceeds approximately 1.7
- The correlation between z and the unobserved unit effect is not too high

Value

A data frame with columns:

variable	Variable name
sd_between	Between-panel standard deviation
sd_within	Within-panel standard deviation
bw_ratio	Ratio of between to within SD

References

Plumper, T., & Troeger, V. E. (2007). Efficient Estimation of Time-Invariant and Rarely Changing Variables in Finite Sample Panel Analyses with Unit Fixed Effects. *Political Analysis*, 15(2), 124-139. doi:10.1093/pan/mpm002

Examples

```
# Create example data
set.seed(42)
N <- 50
T <- 5
id <- rep(1:N, each = T)
z_invariant <- rep(rnorm(N), each = T) # Truly time-invariant
z_slow <- rep(rnorm(N), each = T) + rnorm(N * T, sd = 0.1) # Slowly varying
x_varying <- rnorm(N * T) # Time-varying

data <- data.frame(id = id, z_inv = z_invariant,
                  z_slow = z_slow, x = x_varying)

bw_ratio(data, c("z_inv", "z_slow", "x"), id = "id")
```

xtfifevd

Panel Fixed Effects Estimation for Time-Invariant Variables

Description

Estimates panel models with time-invariant regressors using FEVD, FEF, or FEF-IV methods. Standard fixed effects estimation cannot identify coefficients on time-invariant variables; these methods decompose or filter the unit effects to recover these coefficients.

Usage

```
xtfifevd(
  formula,
  data,
  id,
  time,
  method = c("fevd", "fef", "fef_iv"),
  instruments = NULL,
  na.action = na.omit
)

fevd(formula, data, id, time, na.action = na.omit)

fef(formula, data, id, time, na.action = na.omit)

fef_iv(formula, data, id, time, instruments, na.action = na.omit)
```

Arguments

formula	A formula of the form $y \sim x_1 + x_2 \mid z_1 + z_2$ where variables before \mid are time-varying and variables after \mid are time-invariant.
data	A data frame containing the variables.
id	Character string naming the panel (individual) identifier variable.
time	Character string naming the time identifier variable.
method	Estimation method: "fevd" (default), "fef", or "fef_iv".
instruments	For method = "fef_iv", a one-sided formula specifying instrumental variables, e.g., $\sim iv_1 + iv_2$.
na.action	How to handle missing values. Default is na.omit.

Details

Model:

The panel model is:

$$y_{it} = \alpha_i + x'_{it}\beta + z'_i\gamma + \varepsilon_{it}$$

where x_{it} are time-varying regressors, z_i are time-invariant regressors, and α_i are individual fixed effects.

Stage 1 (All Methods):

Fixed effects regression of y_{it} on x_{it} yields consistent $\hat{\beta}$ and combined residuals $\hat{u}_{it} = \hat{\alpha}_i + \hat{\varepsilon}_{it}$.

Stage 2:

Time-averaged residuals \bar{u}_i are regressed on z_i :

- **FEF**: OLS of \bar{u}_i on z_i with intercept
- **FEF-IV**: 2SLS using instruments r_i
- **FEVD**: Same as FEF, then Stage 3 pooled OLS (point estimates identical)

Variance Estimation:

Uses Pesaran-Zhou (2016) Equation 17/51 which properly accounts for estimation uncertainty from Stage 1. The naive pooled OLS standard errors from FEVD Stage 3 are **inconsistent** and can understate true SEs by factors of 2-5x or more.

Value

An object of class "xtfffevd" containing:

coefficients Named vector of all coefficients (beta, gamma, intercept)

vcov Variance-covariance matrix using Pesaran-Zhou estimator

beta Coefficients on time-varying variables (from FE stage)

gamma Coefficients on time-invariant variables

intercept Overall intercept

residuals Idiosyncratic residuals from FE stage

fitted.values Fitted values

sigma2_e Variance of idiosyncratic error
sigma2_u Variance of unit effects
N Total number of observations
N_g Number of groups (panels)
T_bar Average time periods per panel
method Estimation method used
call The matched call

Functions

- fevd: FEVD estimation (3-stage, Plümper-Troeger)
- fef: FEF estimation (2-stage, Pesaran-Zhou)
- fef_iv: FEF-IV estimation with instruments

References

Plumper, T., & Troeger, V. E. (2007). Efficient Estimation of Time-Invariant and Rarely Changing Variables in Finite Sample Panel Analyses with Unit Fixed Effects. *Political Analysis*, 15(2), 124-139. doi:10.1093/pan/mpm002

Pesaran, M. H., & Zhou, Q. (2018). Estimation of time-invariant effects in static panel data models. *Econometric Reviews*, 37(10), 1137-1171. doi:10.1080/07474938.2016.1222225

See Also

[fevd](#), [fef](#), [fef_iv](#), [bw_ratio](#)

Examples

```
# Simulate panel data
set.seed(123)
N <- 100 # panels
T <- 10 # time periods
n <- N * T

# Generate data
id <- rep(1:N, each = T)
time <- rep(1:T, N)
alpha_i <- rep(rnorm(N), each = T) # Fixed effects
z <- rep(rnorm(N), each = T) # Time-invariant
x <- rnorm(n) # Time-varying
y <- 1 + 2 * x + 0.5 * z + alpha_i + rnorm(n, sd = 0.5)

data <- data.frame(id = id, time = time, y = y, x = x, z = z)

# Estimate with different methods
fit_fevd <- xtfifevd(y ~ x | z, data = data, id = "id", time = "time")
summary(fit_fevd)
```

```
fit_fef <- xtfifevd(y ~ x | z, data = data, id = "id", time = "time",  
                  method = "fef")  
summary(fit_fef)
```

Index

`bw_ratio`, [2](#), [5](#)

`fef`, [5](#)

`fef(xtfifevd)`, [3](#)

`fef_iv`, [5](#)

`fef_iv(xtfifevd)`, [3](#)

`fevd`, [5](#)

`fevd(xtfifevd)`, [3](#)

`xtfifevd`, [3](#)